

1 **Programmed DNA elimination of germline development genes in songbirds**

2

3 Cormac M. Kinsella^{1,2*}, Francisco J. Ruiz-Ruano^{3*}, Anne-Marie Dion-Côté^{1,4}, Alexander J.
4 Charles⁵, Toni I. Gossmann⁵, Josefa Cabrero³, Dennis Kappei⁶, Nicola Hemmings⁵, Mirre J. P.
5 Simons⁵, Juan P. M. Camacho³, Wolfgang Forstmeier⁷, Alexander Suh¹

6

7 ¹Department of Evolutionary Biology, Evolutionary Biology Centre (EBC), Science for Life
8 Laboratory, Uppsala University, SE-752 36, Uppsala, Sweden.

9 ²Laboratory of Experimental Virology, Department of Medical Microbiology, Amsterdam UMC,
10 University of Amsterdam, 1105 AZ, Amsterdam, The Netherlands.

11 ³Department of Genetics, University of Granada, E-18071, Granada, Spain.

12 ⁴Department of Molecular Biology & Genetics, Cornell University, NY 14853, Ithaca, United
13 States.

14 ⁵Department of Animal and Plant Sciences, University of Sheffield, S10 2TN, Sheffield, United
15 Kingdom.

16 ⁶Cancer Science Institute of Singapore, National University of Singapore, 117599, Singapore.

17 ⁷Max Planck Institute for Ornithology, D-82319, Seewiesen, Germany.

18

19 *These authors contributed equally to this work (alphabetical order).

20

21 Correspondence and requests for materials should be addressed to F.J.R.R. (email:
22 fjruizruano@ugr.es) and A.S. (email: alexander.suh@ebc.uu.se).

23

24 **Summary**

25 Genomes can vary within individual organisms. Programmed DNA elimination leads to dramatic
26 changes in genome organisation during the germline–soma differentiation of ciliates¹, lampreys²,
27 nematodes^{3,4}, and various other eukaryotes⁵. A particularly remarkable example of tissue-specific
28 genome differentiation is the germline-restricted chromosome (GRC) in the zebra finch which is
29 consistently absent from somatic cells⁶. Although the zebra finch is an important animal model
30 system⁷, molecular evidence from its large GRC (>150 megabases) is limited to a short intergenic
31 region⁸ and a single mRNA⁹. Here, we combined cytogenetic, genomic, transcriptomic, and
32 proteomic evidence to resolve the evolutionary origin and functional significance of the GRC.
33 First, by generating tissue-specific *de-novo* linked-read genome assemblies and re-sequencing
34 two additional germline and soma samples, we found that the GRC contains at least 115 genes
35 which are paralogous to single-copy genes on 18 autosomes and the Z chromosome. We detected
36 an amplification of ≥ 38 GRC-linked genes into high copy numbers (up to 308 copies) but,
37 surprisingly, no enrichment of transposable elements on the GRC. Second, transcriptome and
38 proteome data provided evidence for functional expression of GRC genes at the RNA and protein
39 levels in testes and ovaries. Interestingly, the GRC is enriched for genes with highly expressed
40 orthologs in chicken gonads and gene ontologies involved in female gonad development. Third,
41 we detected evolutionary strata of GRC-linked genes. Developmental genes such as *bicc1* and
42 *trim71* have resided on the GRC for tens of millions of years, whereas dozens have become
43 GRC-linked very recently. The GRC is thus likely widespread in songbirds (half of all bird
44 species) and its rapid evolution may have contributed to their diversification. Together, our
45 results demonstrate a highly dynamic evolutionary history of the songbird GRC leading to
46 dramatic germline–soma genome differences as a novel mechanism to minimise genetic conflict
47 between germline and soma.

48 **Text**

49 Not all cells of an organism must contain the same genome. Some eukaryotes exhibit dramatic
50 differences between their germline and somatic genomes, resulting from programmed DNA
51 elimination of chromosomes or fragments thereof during germline–soma differentiation⁵. Here
52 we present the first comprehensive analyses of a germline-restricted chromosome (GRC). The
53 zebra finch (*Taeniopygia guttata*) GRC is the largest chromosome of this songbird⁶ and likely
54 comprises >10% of the genome (>150 megabases)^{7,10}. Cytogenetic evidence suggests the GRC is
55 inherited through the female germline, expelled late during spermatogenesis, and eliminated from
56 the soma during early embryo development^{6,11}. Previous analyses of a 19-kb intergenic region
57 suggested that the GRC contains sequences with high similarity to regular chromosomes (‘A
58 chromosomes’)⁸.

59

60 In order to reliably identify sequences as GRC-linked, we used a single-molecule sequencing
61 technology not applied previously in birds that permits reconstruction of long haplotypes through
62 linked reads¹². We generated separate haplotype-resolved *de-novo* genome assemblies for the
63 germline and soma of a male zebra finch (testis and liver; ‘Seewiesen’; Supplementary Table 1).
64 We further used the linked-read data to compare read coverage and haplotype barcode data in
65 relation to the zebra finch somatic reference genome (‘taeGut2’)⁷, allowing us to identify
66 sequences that are shared, amplified, or unique to the germline genome in a fashion similar to
67 recent studies on cancer aneuploidies¹³. We also re-sequenced the germline and soma from two
68 unrelated male zebra finches (‘Spain’; testis and muscle; Extended Data Fig. 1) using short reads.

69

70 We first established the presence of the GRC in the three germline samples. Cytogenetic analysis
71 using fluorescence *in-situ* hybridisation (FISH) with a new GRC probe showed that the GRC is
72 present exclusively in the germline and eliminated during spermatogenesis as hypothesised (Fig.
73 1a-b, Extended Data Fig. 2)^{6,11}. We compared germline/soma sequencing coverage by mapping
74 reads from all three sampled zebra finches onto the reference genome assembly (regular ‘A
75 chromosomes’), revealing consistently germline-increased coverage for single-copy regions,
76 reminiscent of programmed DNA elimination of short genome fragments in lampreys² (Fig. 1c-d).
77 A total of 92 regions (41 with >10 kb length) on 13 chromosomes exhibit >4-fold increased
78 germline coverage in ‘Seewiesen’ relative to the soma (Fig. 1e, Supplementary Table 2). Such a
79 conservative coverage cut-off provides high confidence in true GRC-amplified regions. We
80 obtained nearly identical confirmatory results using another library preparation method for the
81 ‘Spain’ birds (Fig. 1f). Notably, the largest block of testis-increased coverage spans nearly 1 Mb
82 on chromosome 1 and overlaps with the previously⁸ FISH-verified intergenic region 27L4 (Fig.
83 1e-f).

84
85 Our linked-read and re-sequencing approach allowed us to determine the sequence content of the
86 GRC. The GRC is effectively a non-recombining chromosome as it recombines with itself after
87 duplication, probably to ensure stable inheritance during female meiosis⁸. We predicted that the
88 GRC would be highly enriched in repetitive elements, similar to the female-specific avian W
89 chromosome (repeat density >50%, compared to <10% genome-wide)¹⁴. Surprisingly, neither
90 assembly-based nor read-based repeat quantifications detected a significant enrichment in
91 transposable elements or satellite repeats in the germline samples relative to the soma samples
92 (Extended Data Figure 3, Supplementary Table 3). Instead, most germline coverage peaks lie in

93 single-copy regions of the reference genome overlapping 38 genes (Fig. 1e-f, Table 1,
94 Supplementary Table 4), suggesting that these peaks stem from very similar GRC-amplified
95 paralogs with high copy numbers (up to 308 copies per gene; Supplementary Table 5). GRC
96 linkage of these regions is further supported by sharing of linked-read barcodes between different
97 amplified chromosomal regions in germline but not soma (Fig. 1g-h), suggesting that these
98 regions reside on the same haplotype (Extended Data Fig. 4). We additionally identified 245
99 GRC-linked genes through germline-specific single-nucleotide variants (SNVs) present in read
100 mapping of all three germline samples onto zebra finch reference genes (up to 402 SNVs per
101 gene; Supplementary Table 4). As a control, we used the same methodology to screen for soma-
102 specific SNVs and found no such genes. We conservatively consider the 38 GRC-amplified
103 genes and those with at least 5 germline-specific SNVs as our highest-confidence set (Table 1).
104 We also identified GRC-linked genes using germline–soma assembly subtraction (Fig. 1i);
105 however, all were already found via coverage or SNV evidence (Table 1). Together with the *napa*
106 gene recently identified in transcriptomes (Fig. 1j)⁹, our complementary approaches yielded 115
107 high-confidence GRC-linked genes with paralogs located on 18 autosomes and the Z
108 chromosome (Table 1; all 267 GRC genes in Supplementary Table 4).

109
110 We next tested whether the GRC is functional and thus probably physiologically important using
111 transcriptomics and proteomics. We sequenced RNA from the same tissues of the two Spanish
112 birds used for genome re-sequencing and combined these with published testis and ovary RNA-
113 seq data from North American domesticated zebra finches^{9,15}. Among the 115 high-confidence
114 genes, 6 and 32 were transcribed in testes and ovaries, respectively (Table 1). Note, these are
115 only genes for which we could reliably separate GRC-linked and A-chromosomal paralogs using

116 GRC-specific SNVs in the transcripts (Fig. 2a-b, Extended Data Fig. 5, Supplementary Table 6).
117 We next verified translation of GRC-linked genes through protein mass spectrometry data for 7
118 testes and 2 ovaries from another population ('Sheffield'). From 83 genes with GRC-specific
119 amino acid changes, we identified peptides from 5 GRC-linked genes in testes and ovaries (Fig.
120 2c-d, Extended Data Fig. 6, Table 1). We therefore established that many GRC-linked genes are
121 transcribed and translated in adult male and female gonads, extending previous RNA evidence
122 for a single gene⁹ and questioning the hypothesis from cytogenetic studies that the GRC is
123 silenced in the male germline^{16,17}. Instead, we propose that the GRC has important functions
124 during germline development, which is supported by a significant enrichment in gene ontology
125 terms related to reproductive developmental processes among GRC-linked genes (Fig. 2e,
126 Supplementary Table 7). We further found that the GRC is significantly enriched in genes that
127 are also germline-expressed in GRC-lacking species with RNA expression data available from
128 many tissues¹⁸ (Fig. 2f, Supplementary Table 8). Specifically, out of 65 chicken orthologs of
129 high-confidence GRC-linked genes, 22 and 6 are most strongly expressed in chicken testis and
130 ovary, respectively.

131
132 The observation that all identified GRC-linked genes have A-chromosomal paralogs allowed us
133 to decipher the evolutionary origins of the GRC. We utilised phylogenies of GRC-linked genes
134 and their A-chromosomal paralogs to infer when these genes copied to the GRC, similarly to the
135 inference of evolutionary strata of sex chromosome differentiation¹⁹. First, the phylogeny of the
136 intergenic 27L4 locus of our germline samples and a previous GRC sequence⁸ demonstrated
137 stable inheritance among the sampled zebra finch populations (Fig. 3a). Second, 37 gene trees of
138 GRC-linked genes with germline-specific SNVs and available somatic genome data from other

139 birds identify at least five evolutionary strata (Fig. 3b-f, Extended Data Fig. 7, Table 1), with all
140 but stratum 3 containing expressed genes (*cf.* Fig. 2a-d). Stratum 1 emerged during early songbird
141 diversification, stratum 2 before the diversification of estrildid finches, and stratum 3 within
142 estrildid finches (Fig. 3g). The presence of at least 7 genes in these three strata implies that the
143 GRC is tens of millions of years old and likely present across songbirds (Extended Data Fig. 7),
144 consistent with a recent cytogenetics preprint²⁰. Notably, stratum 4 is specific to the zebra finch
145 species and stratum 5 to the Australian zebra finch subspecies (Fig. 3g), suggesting piecemeal
146 addition of genes from 18 autosomes and the Z chromosome over millions of years of GRC
147 evolution (Fig. 3h). The long-term residence of expressed genes on the GRC implies that they
148 have been under selection, such as *bicc1* and *trim71* on GRC stratum 1 whose human orthologs
149 are important for embryonic cell differentiation²¹. Using ratios of non-synonymous to
150 synonymous substitutions (dN/dS) for GRC-linked genes with >50 GRC-specific SNVs, we
151 found 17 genes evolving faster than their A-chromosomal paralogs (Supplementary Table 9).
152 However, we also detected long-term purifying selection on 9 GRC-linked genes, including *bicc1*
153 and *trim71*, as well as evidence for positive selection on *puf60*, again implying that the GRC is an
154 important chromosome with a long evolutionary history.

155
156 Here we provided the first evidence for the origin and functional significance of a GRC. Notably,
157 our analyses suggest that the GRC emerged during early songbird evolution and we predict it to
158 be present in half of all bird species. The species-specific addition of dozens of genes on stratum
159 5 implies that the rapidly evolving GRC likely contributed to reproductive isolation during the
160 massive diversification of songbirds²². It was previously hypothesised that GRCs are formerly
161 parasitic B chromosomes that became stably inherited^{23,24}. Our evidence for an enrichment of

162 germline-expressed genes on the zebra finch GRC is reminiscent of nematodes and lampreys
163 where short genome fragments containing similar genes are eliminated during germline–soma
164 differentiation²⁻⁴. All these cases constitute extreme mechanisms of gene regulation through
165 germline–soma gene removal rather than transcriptional repression^{3,5,10}. Remarkably, the GRC
166 harbours several genes involved in the control of cell division and germline determination,
167 including *prdm1*, a key regulator of primordial germ cell differentiation in mice^{25,26}.
168 Consequently, we hypothesise that the GRC became indispensable for its host by the acquisition
169 of germline development genes and probably acts as a germline-determining chromosome. The
170 aggregation of developmental genes on a single eliminated chromosome constitutes a novel
171 mechanism to ensure germline-specific gene expression in multicellular organisms. This may
172 allow adaptation to germline-specific functions free of detrimental effects on the soma which
173 would otherwise arise from antagonistic pleiotropy.

174

175 **References (max. 30 references)**

- 176 1 Chen, X. *et al.* The architecture of a scrambled genome reveals massive levels of genomic
177 rearrangement during development. *Cell* **158**, 1187-1198 (2014).
- 178 2 Smith, J. J. *et al.* The sea lamprey germline genome provides insights into programmed
179 genome rearrangement and vertebrate evolution. *Nat. Genet.* **50**, 270-277 (2018).
- 180 3 Wang, J. *et al.* Silencing of germline-expressed genes by DNA elimination in somatic
181 cells. *Dev. Cell* **23**, 1072-1080 (2012).
- 182 4 Wang, J. *et al.* Comparative genome analysis of programmed DNA elimination in
183 nematodes. *Genome Res.* **27**, 2001-2014 (2017).
- 184 5 Wang, J. & Davis, R. E. Programmed DNA elimination in multicellular organisms. *Curr.*
185 *Opin. Genet. Dev.* **27**, 26-34 (2014).

- 186 6 Pigozzi, M. I. & Solari, A. J. Germ cell restriction and regular transmission of an
187 accessory chromosome that mimics a sex body in the zebra finch, *Taeniopygia guttata*.
188 *Chromosome Res.* **6**, 105-113 (1998).
- 189 7 Warren, W. C. *et al.* The genome of a songbird. *Nature* **464**, 757–762 (2010).
- 190 8 Itoh, Y., Kampf, K., Pigozzi, M. I. & Arnold, A. P. Molecular cloning and
191 characterization of the germline-restricted chromosome sequence in the zebra finch.
192 *Chromosoma* **118**, 527-536 (2009).
- 193 9 Biederman, M. K. *et al.* Discovery of the first germline-restricted gene by subtractive
194 transcriptomic analysis in the zebra finch, *Taeniopygia guttata*. *Curr. Biol.* **28**, 1620-1627
195 (2018).
- 196 10 Smith, J. J. Programmed DNA elimination: keeping germline genes in their place. *Curr.*
197 *Biol.* **28**, R601-R603 (2018).
- 198 11 Pigozzi, M. I. & Solari, A. J. The germ-line-restricted chromosome in the zebra finch:
199 recombination in females and elimination in males. *Chromosoma* **114**, 403-409 (2005).
- 200 12 Weisenfeld, N. I., Kumar, V., Shah, P., Church, D. M. & Jaffe, D. B. Direct determination
201 of diploid genome sequences. *Genome Res.* **27**, 757-767 (2017).
- 202 13 Bell, J. M. *et al.* Chromosome-scale mega-haplotypes enable digital karyotyping of cancer
203 aneuploidy. *Nucleic Acids Res.* **45**, e162-e162 (2017).
- 204 14 Kapusta, A. & Suh, A. Evolution of bird genomes—a transposon's-eye view. *Ann. N. Y.*
205 *Acad. Sci.* **1389**, 164–185 (2017).
- 206 15 Singhal, S. *et al.* Stable recombination hotspots in birds. *Science* **350**, 928-932 (2015).
- 207 16 del Priore, L. & Pigozzi, M. I. Histone modifications related to chromosome silencing and
208 elimination during male meiosis in Bengalese finch. *Chromosoma* **123**, 293-302 (2014).
- 209 17 Goday, C. & Pigozzi, M. I. Heterochromatin and histone modifications in the germline-
210 restricted chromosome of the zebra finch undergoing elimination during spermatogenesis.
211 *Chromosoma* **119**, 325-336 (2010).
- 212 18 Marin, R. *et al.* Convergent origination of a *Drosophila*-like dosage compensation
213 mechanism in a reptile lineage. *Genome Res.* **27**, 1974-1987 (2017).
- 214 19 Lahn, B. T. & Page, D. C. Four evolutionary strata on the human X chromosome. *Science*
215 **286**, 964-967 (1999).
- 216 20 Torgasheva, A. A. *et al.* Germline-restricted chromosome (GRC) is widespread among
217 songbirds. *bioRxiv* doi:10.1101/414276 (2018).

- 218 21 Uhlén, M. *et al.* Tissue-based map of the human proteome. *Science* **347** (2015).
- 219 22 Moyle, R. G. *et al.* Tectonic collision and uplift of Wallacea triggered the global songbird
220 radiation. *Nat. Commun.* **7**, 12709 (2016).
- 221 23 Camacho, J. P. M. B chromosomes. in *The Evolution of the Genome* (ed T. Ryan
222 Gregory) 223-286 (Elsevier Academic Press, 2005).
- 223 24 Camacho, J. P. M., Sharbel, T. F. & Beukeboom, L. W. B-chromosome evolution. *Philos.*
224 *Trans. R. Soc. B* **355**, 163-178 (2000).
- 225 25 Ohinata, Y. *et al.* Blimp1 is a critical determinant of the germ cell lineage in mice. *Nature*
226 **436**, 207 (2005).
- 227 26 Vincent, S. D. *et al.* The zinc finger transcriptional repressor Blimp1/Prdm1 is
228 dispensable for early axis formation but is required for specification of primordial germ
229 cells in the mouse. *Development* **132**, 1315-1325 (2005).
- 230 27 Hooper, D. M. & Price, T. D. Rates of karyotypic evolution in Estrildid finches differ
231 between island and continental clades. *Evolution* **69**, 890-903 (2015).
- 232 28 Mossman, J. A., Birkhead, T. R. & Slate, J. O. N. The whole mitochondrial genome
233 sequence of the zebra finch (*Taeniopygia guttata*). *Molecular Ecology Notes* **6**, 1222-
234 1227 (2006).

235

236 **Acknowledgements** We thank Peter Ellis, Moritz Hertel, Martin Irestedt, Regine Jahn, Max
237 Käller, Bart Kempnaers, Ulrich Knief, Pedro Lanzas, Juan Gabriel Martínez, Julio Mendo-
238 Hernández, Beatriz Navarro-Domínguez, Remi-André Olsen, Mattias Ormestad, Yifan Pei,
239 Douglas Scofield, Linnéa Smeds, Venkat Talla, and members of the Barbash lab and the Suh lab
240 for support and discussions. Mozes Blom, Jesper Boman, Nazeefa Fatima, James Galbraith,
241 Octavio Palacios, and Matthias Weissensteiner provided helpful comments on an earlier version
242 of this manuscript. A.S. was supported by grants from the Swedish Research Council Formas
243 (2017-01597), the Swedish Research Council Vetenskapsrådet (2016-05139), and the SciLifeLab
244 Swedish Biodiversity Program (2015-R14). The Swedish Biodiversity Program has been made
245 available by support from the Knut and Alice Wallenberg Foundation. F.J.R.R., J.C., and J.P.M.C.
246 were supported by the Spanish Secretaría de Estado de Investigación, Desarrollo e Innovación
247 (CGL2015-70750-P), including FEDER funds, and F.J.R.R. was also supported by a Junta de
248 Andalucía fellowship. A.M.D.C was supported by a postdoc fellowship from Sven och Lilly

249 Lawskis fond. T.I.G. was supported by a Leverhulme Early Career Fellowship Grant (ECF-2015-
250 453). T.I.G., A.J.C. (CABM DTP), and M.S. were supported by a NERC grant (NE/N013832/1).
251 N.H. was supported by a Patrick & Irwin-Packington Fellowship from the University of Sheffield
252 and a Royal Society Dorothy Hodgkin Fellowship. D.K. was supported by the National Research
253 Foundation Singapore and the Singapore Ministry of Education under its Research Centres of
254 Excellence initiative. W.F. was supported by the Max Planck Society. Some of the computations
255 were performed on resources provided by the Swedish National Infrastructure for Computing
256 (SNIC) through Uppsala Multidisciplinary Center for Advanced Computational Science
257 (UPPMAX). The authors acknowledge support from the National Genomics Infrastructure in
258 Stockholm funded by Science for Life Laboratory, the Knut and Alice Wallenberg Foundation
259 and the Swedish Research Council.

260
261 **Author Contributions** Conceptualisation: W.F., A.S., J.P.M.C., F.J.R.R., C.M.K., A.M.D.C.,
262 T.I.G.; cytogenetics analyses and interpretation: J.P.M.C., F.J.R.R., J.C.; genomic analyses and
263 interpretation: A.S., C.M.K., F.J.R.R., A.M.D.C.; transcriptomic analyses and interpretation:
264 F.J.R.R.; proteomic analyses and interpretation: T.I.G., A.J.C., D.K., M.J.P.S., N.H.; gene
265 enrichment analyses and interpretation: C.M.K., W.F., A.S.; phylogenetic analyses and
266 interpretation: F.J.R.R., A.S., C.M.K., T.I.G.; manuscript writing: A.S. with input from all
267 authors; methods and supplements writing: C.M.K. with input from all authors; supervision: A.S.,
268 J.P.M.C., T.I.G., M.J.P.S. All authors read and approved the manuscript.

269
270 **Author Information** The authors declare no competing financial interests. Correspondence and
271 requests for materials should be addressed to F.J.R.R. (email: fjuizruano@ugr.es) and A.S.
272 (alexander.suh@ebc.uu.se).

273
274 **Tables and Figures**

275

276 **Table 1 | The 115 high-confidence genes on the GRC with information on their A-chromosomal origin in the**
 277 **reference genome taeGut2, number of testis-specific SNVs, methods supporting their GRC linkage,**
 278 **testis/ovary RNA expression of the GRC paralog, testis/ovary protein expression of the GRC paralog, and**
 279 **evolutionary stratum on the GRC.**

Gene symbol	Chr.	Start	End	SNVs	Method	RNA evidence	Protein evidence	GRC stratum
AAGAB	10	19608548	19634367	10	SNVs			S5
ADGRL2	8	14047115	14171612	10	SNVs			
ADGRL3	4	14919933	15404594	8	SNVs	ovary		
AKIRIN2	3	78683482	78688947	6	SNVs	ovary		S5
ALDH18A1	6	36280145	36301392	17	SNVs			S4
ALG13	4A	18474239	18501426	19	SNVs	ovary		
ARMC6	28	4942046	4946063	5	SNVs			
ATP2A2	15	2841010	2879975	8	SNVs			
BICC1	6	6355408	6434911	402	SNVs	ovary		S1
BMP15	4A	15596686	15598225	29	SNVs, coverage	ovary		S5
BMPR1B	4	18997710	19024248	47	SNVs, coverage			S5
CCND3	26_random			14	SNVs			
CD164	3	69169111	69174605	38	SNVs, coverage	ovary		
COPS2	10	10200701	10222248	1	SNVs, coverage	ovary		
CPEB1	10	3114181	3137661	114	SNVs	ovary		
CSNK1A1L	Un	135422201	135425792	NA	coverage			
CXCL14	13	9423543	9433139	12	SNVs			S5
DDX49	28	4913058	4918451	5	SNVs	ovary		
DIS3L	10	19097281	19112154	13	SNVs	ovary		S5
DNAAF5	14	13758049	13780402	NA	coverage			
DNAH5	2	81235805	81361091	7	SNVs			
DPH6	5	31543945	31606965	13	SNVs, coverage			
EFNB1	4A	5764021	5807953	86	SNVs	ovary		S5
ELAVL4	8	21034240	21098310	364	SNVs	ovary		

EPPK1	Un			52	SNVs			
FBXO16	3	112541865	112568948	6	SNVs			
FEM1B	10	19886491	19891616	9	SNVs	ovary		S5
FIG4	3	69023384	69073678	17	SNVs			S5
FRS3	26_random			42	SNVs, coverage			S5
GBE1	1	105820640	105934310	4	SNVs, coverage			
INTS9	3	112259951	112313512	NA	coverage			
LIAS	4	48132714	48139736	42	SNVs			S2
LIN54	4	13615974	13637371	17	SNVs			
LINC02027	1	106086596	106087033	NA	coverage			
LMBRD2	Z	41646446	41665840	NA	coverage			
LOC100223190	Z	69149414	69156994	41	SNVs			
LOC100224235	Un			5	SNVs			S5
LOC100225322	1A	47543094	47544622	6	SNVs	ovary		
LOC100227189	Un	150797142	150801997	NA	coverage			
LOC100228170	Un	55540047	55541360	NA	coverage			
LOC101233087	Z	47991391	47994344	7	SNVs			
LOC101233688	5	937818	939059	5	SNVs			S5
LOC101233767	18	8034939	8038005	11	SNVs			
LOC101233800	Un			16	SNVs			
LOC101234253	10	19184028	19186114	7	SNVs	ovary		S5
LOC105758464	23	46808	60360	14	SNVs			S5
LOC105758894	26_random			5	SNVs			


LOC10575897 6	2	34301994	34306899	16	SNVs			
LOC10575910 1	3	76396180	76401262	21	SNVs			
LOC10575916 7	4A	15573874	15574621	5	SNVs			
LOC10575919 5	4	14453003	14473747	18	SNVs			
LOC10575919 9	4	20714525	20720872	11	SNVs			
LOC10575926 0	5	1874731	1886007	32	SNVs			S5
LOC10575964 6	Un			7	SNVs			
LOC10575965 5	Un			8	SNVs			
LOC10575966 0	Un			18	SNVs			
LOC10575966 5	Un			5	SNVs			
LOC10575969 2	Un			12	SNVs			
LOC10575991 9	Un			8	SNVs			
LOC10576001 1	Un			7	SNVs			
LOC10576012 3	Un			18	SNVs			
LOC10576022 8	Un			14	SNVs			
LOC10576028 6	Un			18	SNVs			
LOC10576046 1	Un			10	SNVs			
LOC10576087 4	Z	60949696	60953194	19	SNVs	testis		
LOC10576093 6	16_rando m			12	SNVs			
LUC7L3	Un	35019850	35021569	NA	coverage			
MED20	26_rando m	110500	113183	28	SNVs, coverage			S5

MSH4	8	27964612	27983306	30	SNVs			S4
NAPA	NA			NA	Biederman et al. 2018		both	
NEUROG1	13	9450787	9451086	6	SNVs			
NFYA	26	4725655	4735626	7	SNVs			S5
NRBP2	2	156379345	156398225	48	SNVs			
PCSK4	28	4059367	4063775	21	SNVs			
PGC	26_random			24	SNVs			
PHKA1	4A	15562688	15593666	16	SNVs			
PIM1	26	603349	607242	50	SNVs	testis		
PIM3	1A	18426716	18430551	81	SNVs	ovary		
PMM1	1A	49038672	49047011	NA	coverage			
PRDM1	3	70624594	70644625	12	SNVs			
PRKAR1A	18	2200317	2211579	NA	coverage			
PRKAR1B	14	13784578	13872733	NA	coverage			
PRPSAP1	18	8008870	8033058	7	SNVs, coverage	ovary		S5
PSIP1	Z	59887174	59919902	57	SNVs, coverage	ovary		S3
PUF60	2	156354670	156376091	63	SNVs	ovary		
RFC1	4	48169638	48202709	77	SNVs	ovary		S2
RNF157	18	8048721	8062403	NA	coverage			
RNF17	1	45827734	45870640	69	SNVs	ovary	testis	S4
RNF20	Z_random			9	SNVs, subtraction	both		
ROBO1	1	107094521	107228509	19	SNVs, coverage	ovary		S5
ROBO2	1	107529365	107979302	25	SNVs			
RXRA	17	8320685	8355067	14	SNVs			S5
SCRIB	2	156239884	156325797	83	SNVs	ovary		S5
SECISBP2L	10	10159176	10193647	60	SNVs, coverage	ovary	both	S5


SHC4	10	10124441	10151124	11	SNVs, coverage			S4
SPHK1	18	7991834	7994408	2	SNVs, coverage	testis		
SRRT	Un			16	SNVs	both		
SUGP2	28	4930094	4937971	33	SNVs	ovary	both	S5
SURF4	17	7682661	7693000	50	SNVs	ovary		S3
TFEB	26_random	20475	21840	11	SNVs			S5
TIAM2	3	54800961	54890499	NA	coverage			
TRIM71	2	60893878	60907039	159	SNVs, subtraction			S1
UBE2O	18	7960889	7981633	NA	coverage			
UGDH	4	48113314	48126079	136	SNVs, coverage, subtraction	ovary	ovary	S2
UNC5C	4	19035187	19126466	13	SNVs, coverage			
Unnamed	Un	124574513	124575553	NA	coverage			
Unnamed	Un	127129819	127130503	NA	coverage			
Unnamed	16_random	26580	73126	NA	coverage			
Unnamed	Un	130103514	130104264	NA	coverage			
Unnamed	Un	50859565	50860210	NA	coverage			
Unnamed	Un	115355883	115358154	NA	coverage			
Unnamed	Un	124578595	124579326	NA	coverage			
VEGFA	3	31631385	31652650	34	SNVs, coverage	both		
WDR19	4	48204115	48240398	34	SNVs	ovary		S5
ZWILCH	10	19199771	19206407	8	SNVs	ovary		S5

280 Note: We were able to place only some genes on evolutionary strata due to our strict criteria for evaluating the
 281 maximum likelihood gene trees. The remaining genes lacked sequence information from several of the other sampled
 282 somatic genomes or had poorly resolved tree topologies.

283




284
 285 **Figure 1 | The zebra finch germline-restricted chromosome contains genes copied from many A chromosomes.**
 286 **a-b**, Cytogenetic evidence for GRC absence in muscle (**a**) and GRC presence in the testis (**b**) of the same bird
 287 (Spain_1) using fluorescence *in-situ* hybridisation (FISH) of our new GRC-ampliconic probe *dph6* (selected due its
 288 high germline/soma coverage ratio; cf. panels e-f). The scale bar indicates 10 μ m. **c-d**, Comparison of germline/soma
 289 coverage ratios (red) for 1 kb windows with an expected symmetrical distribution (blue) indicates enrichment of
 290 single-copy regions in the germline, similar to lamprey² both in Spain (**c**; average of Spain_1 and Spain_2 coverage;
 291 PCR-free short reads) and Seewiesen (**d**; linked reads) samples. Y-axis is truncated for visualisation. **e-f**, Manhattan
 292 plot of germline/soma coverage ratios in 1 kb windows across chromosomes of the somatic reference genome
 293 *taeGut2*. Colours indicate high-confidence GRC-linked genes and their identification (red: coverage, blue: SNVs,
 294 purple: both; Table 1). Note that the similarities between Seewiesen (**e**) and Spain_1/Spain_2 averages (**f**) constitute
 295 independent biological replicates for GRC-ampliconic regions, as the data are based on different domesticated
 296 populations and different library preparation methods. Red arrows denote two FISH-verified GRC-amplified regions
 297 (cf. panel b)⁸. Only chromosomes >5 Mb are shown for clarity. **g-h**, Linked-read barcode interaction heatmaps of an
 298 inter-chromosomal rearrangement on the GRC absent in Seewiesen liver (**g**) but present in Seewiesen testis (**h**). **i-j**,
 299 Coverage plots of two examples of GRC-linked genes that are divergent from their A-chromosomal paralog, *trim71*
 300 (**i**) and *napa* (**j**)⁹, and thus have very low coverage (normalised by total reads and genome size) in soma.




301
 302 **Figure 2 | The zebra finch germline-restricted chromosome is expressed in male and female gonads.** a-b,
 303 Comparison of coverage and read pileups for DNA-seq data from Spain_1 and Spain_2 testis/muscle, RNA-seq data
 304 from Spain_1 and Spain_2 testis, and available ovary RNA-seq data⁹. Shown are 100-bp regions within *trim71* (a)
 305 and *bicc1* (b). Colours indicate SNVs deviating from the reference genome *taeGut2*. c, Example alignments of
 306 proteomics data showing a subset of peptide expression of the respective GRC-linked paralog of *ugdh* and *napa*
 307 (alternative or ‘alt’; cf. reference or ‘ref’). d, Proteomic evidence for GRC protein expression (‘alt’) in comparison to
 308 their A-chromosomal paralog (‘ref’) of 5 genes in 7 sampled testes and 2 sampled ovaries. For label-free
 309 quantification (LFQ), unique as well as razor (non-unique) peptides were used. Note that unique peptides may occur
 310 in several of the 9 samples. e, Gene ontology term enrichment analysis of the 115 high-confidence GRC-linked genes
 311 (77 mapped gene symbols). Colours indicate the log₁₀ of the false discovery rate-corrected *p*-value, circle sizes
 312 denote fold enrichment above expected values. f, Expression evidence for orthologs of three different sets of GRC
 313 genes in testes, ovaries, or other tissues in chicken¹⁸. Randomisation tests show a significant enrichment for
 314 germline-expressed genes among the 115 high-confidence GRC genes and all 267 GRC genes, but not the 38
 315 ampliconic GRC genes.
 316

317




318

319 **Figure 3 | The zebra finch germline-restricted chromosome is ancient and highly dynamic.** **a**, Phylogeny of the
 320 intergenic 27L4 locus previously sequenced by Itoh et al.⁸ suggests stable inheritance of the GRC paralog
 321 (alternative or ‘alt’ in red; *cf.* reference or ‘ref’) among the sampled zebra finches. **b–f**, Phylogenies of GRC-linked
 322 genes (‘alt’, in red; most selected from expressed genes) diverging from their A-chromosomal paralogs (‘ref’)
 323 before/during early songbird evolution (**b**; *bicc1*, stratum 1; *cf.* Extended Data Fig. 7), during songbird evolution (**c**;
 324 *ugdh*, stratum 2), during estrildid finch evolution (**d**; *psip1*, stratum 3), in the ancestor of the zebra finch species (**e**;
 325 *rnf17*, stratum 4), and in the Australian zebra finch subspecies (**f**; *secisbp2l*; stratum 5). The maximum likelihood
 326 phylogenies in panels a–f (only bootstrap values $\geq 50\%$ shown) include available somatic genome data from estrildid
 327 finches and other songbirds. **g**, Species tree of selected songbirds showing the emergence of evolutionary strata (S1–
 328 S5) on the GRC (red gene names). Molecular dates are based on previous phylogenies^{22,27}. Bird illustrations were
 329 used with permission from Lynx Edicions. **h**, Circos plot indicating A-chromosomal origin of high-confidence GRC-
 330 linked genes from 18 autosomes and the Z chromosome. Note that A-chromosomal paralogs of 37 genes remain
 331 unplaced on chromosomes in the current zebra finch reference genome *taeGut2*.



332
333


334 **Extended Data Figure 1** | Maximum likelihood phylogeny of the five zebra finch mitochondrial haplotypes
335 described by Mossman et al.²⁸ and mitogenomes assembled from all zebra finch Illumina libraries used in this work,
336 comprising both the Australian zebra finch (*Taeniopygia guttata castanotis*) and the Timor zebra finch (*Taeniopygia*
337 *guttata guttata*) subspecies. Note that the three individuals sequenced by us (red colour) and by Biederman et al.⁹
338 belong to different mitochondrial haplotypes.
339



340

341 **Extended Data Figure 2 | FISH analysis in testis cells of the Spain_1 zebra finch individual using the *dph6***
342 **probe (red) counterstained with DAPI (blue).** Note the presence of primary (PS) and secondary (SS)
343 spermatocytes, young spermatids (YS) and maturing spermatids at round (RS) and elongating (ES) stages. Also note
344 that the *dph6* probe hybridises with only part of the GRC chromosome (arrow), and this is apparent in PS at
345 leptotene-zygotene (a), pachytene (b-c, e) and in GRCs which failed to integrate into the main nucleus of SS or YS
346 cells (d), with no FISH signal in somatic cells (SC) indicating GRC absence in somatic structural testis cells (c). The
347 half size of GRC in the SS cell in panel c, compared with that in the PS next to it and that those lying outside nuclei
348 in panel d, indicates that GRC sometimes divides equationally in the first meiotic division (resulting in the half sized
349 GRC body in panel c) but, in most cases, it divides reductionally yielding the large sized GRCs in panel d. Note that
350 RS and ES nuclei in panel e lack FISH signal, indicating GRC absence. All photographs were made at the same
351 magnification, and the scale bar in panel a indicates 10 μm .

352



353

354 **Extended Data Figure 3 | The zebra finch GRC is not enriched in satellites or specific transposable element**

355 **families.** **a**, Comparison of spectra for k-mers shared between or exclusive to genome sequencing data from testis

356 and liver of the Seewiesen sample, showing that the germline is not enriched for exclusive high frequency k-mers,

357 but is conspicuously enriched in high frequency k-mers shared with the soma. **b**, Comparison of simple repeat


358 abundance as assessed by kSeek in the Spanish muscle samples relative to the testis samples. **c**, Same as in panel b,

359 with a focus on low abundance simple repeats. **d-e**, Repeat landscapes based on RepeatMasker analyses showing the

360 main repetitive element families for genome re-sequencing data from muscle (**d**) and testis (**e**) of the combined

361 Spanish samples. **f**, Subtractive repeat landscape obtained by subtracting muscle from testis counts showing a general


362 impoverishment of testis for most of the repetitive elements (negative values) due to the presence of the GRC.



363

364 **Extended Data Figure 4 | Testis-specific linked-read barcode sharing between A chromosomes indicates GRC**
 365 **haplotypes.** Plots show side-by-side comparison of the inter-chromosomal barcode overlap for 200-kb regions for
 366 the liver and testis, respectively (chromosome position scale in Mb). With the exception of the interaction between
 367 chromosome 6 and chromosome 8 (bottom right) showing some background in the liver sample (potentially due to a
 368 shared A-chromosomal rearrangement), all inter-chromosomal structural variants were testis-specific and thus
 369 indicative of being on the same haplotype on the GRC. We exported barcode overlap matrices from the Loupe
 370 browser for testis-specific structural variants called by LongRanger and plotted them in R (v. 3.5.1). We reassigned 0
 371 values to “NA” (shown in white on the plot) and \log_2 -transformed all values. Note that the scale varies across plots.


372



373

374 **Extended Data Figure 5 | Further examples for RNA expression of GRC-linked genes.** Comparison of coverage
375 and read pileups for DNA-seq from Spain_1 and Spain_2 testis/muscle, RNA-seq data from Spain_1 and Spain_2
376 testis, and available ovary RNA-seq data⁹. Shown are 100-bp regions within 10 selected genes. Colours indicate
377 SNVs deviating from the zebra finch reference genome *taeGut2*.

378



379

380 **Extended Data Figure 6 | Proteomic evidence for GRC protein presence in zebra finch testes and ovaries.** The

381 five proteins listed at the top are also shown in Fig. 2d, i.e., those where we could differentiate between peptides

382 from GRC vs. A-chromosomes. GRC paralogs are denoted by the 'alt' suffix, whereas A-chromosomal paralogs are

383 denoted by the 'ref' suffix. Unique sequence coverage corresponds to the peptide coverage percentage of the

384 reference protein sequence. Note that unique peptides may occur in several samples (testes/ovaries). Entries of only


385 one protein identification have sufficient evidence at the peptide level to differentiate between the GRC and A-

386 chromosomal paralogs due to coverage of non-identical regions between the both reference sequences; entries of

387 more than one protein identification contain evidence of presence based solely on identical regions, thus cannot be

388 differentiated at the proteomic level. Entries of only one protein identification without the corresponding 'alt' or 'ref'

389 variant contain evidence that span the non-identical region only, thus the alternate variant need not be called.



390
 391 **Extended Data Figure 7 | Gene trees of GRC-linked genes from stratum 1 and their A-chromosomal paralogs**
 392 **from broad taxon sampling imply GRC emergence in the ancestor of Passeriformes.** **a**, Maximum likelihood
 393 gene tree of *trim71* (partitioned for codon positions) suggesting GRC linkage in the ancestor of Passeriformes. **b**,
 394 Maximum likelihood gene tree of *bicc1* (only 3' UTR) suggesting GRC linkage in the ancestor of oscine songbirds.

395
 396

397 **Supplementary Information**

398 **Methods and Supplementary Text**

399 **Supplementary Table 1 | Assembly metrics of linked-read *de-novo* assemblies generated**
400 **from liver and testis samples of the Seewiesen zebra finch individual.**

401 **Supplementary Table 2 | Position, length, and library source of genomic blocks >10-kb with**
402 **average germline/soma corrected coverage >4, with respect to the zebra finch reference**
403 **genome (taeGut2).**

404 **Supplementary Table 3 | Repeat annotation of the pseudohaploid testis and liver *de-novo***
405 **assemblies from the Seewiesen zebra finch individual.**

406 **Supplementary Table 4 | All 267 genes on the GRC with information on their A-**
407 **chromosomal origin in taeGut2, number of testis-specific SNVs, methods supporting their**
408 **GRC linkage, testis/ovary RNA expression of the GRC paralog, testis/ovary protein**
409 **expression of the GRC paralog, and evolutionary stratum on the GRC.**

410 **Supplementary Table 5 | Copy number estimates for 61 GRC-linked genes with at least 2**
411 **copies on the GRC as estimated from excess coverage in testis.**

412 **Supplementary Table 6 | Transcriptome analyses of GRC-linked genes showing the number**
413 **of 'alt' SNVs per transcript with a minimum of 100 reads and an 'alt'/'ref' SNV mapping**
414 **ratio above 1% in testes and ovary RNA-seq data.**

415 **Supplementary Table 7 | Enriched gene ontology terms for 167 mapped gene symbols from**
416 **all 267 GRC-linked genes, and 77 mapped genes from 115 high confidence genes.**

417 **Supplementary Table 8 | Enrichment analyses of GRC gene orthologs in chicken and**
418 **human RNA-seq data for testes, ovaries, and other tissues.**

419 **Supplementary Table 9 | Codon substitution rate analyses for 17 genes with at least 50**
420 **GRC-specific SNVs.**

421 **Supplementary Data**