

## Comment



**Cite this article:** Wedekind C. 2018 A predicted interaction between odour pleasantness and intensity provides evidence for major histocompatibility complex social signalling in women. *Proc. R. Soc. B* **285**: 20172714.  
<http://dx.doi.org/10.1098/rspb.2017.2714>

Received: 5 December 2017

Accepted: 8 January 2018

**Subject Category:**

Behaviour

**Subject Areas:**

behaviour, genetics, evolution

**Author for correspondence:**

Claus Wedekind

e-mail: [claus.wedekind@unil.ch](mailto:claus.wedekind@unil.ch)

The accompanying reply can be viewed at  
<http://dx.doi.org/10.1098/rspb.2018.0566>.

Electronic supplementary material is available online at <https://doi.org/10.6084/m9.figshare.c.4089986>.

# A predicted interaction between odour pleasantness and intensity provides evidence for major histocompatibility complex social signalling in women

Claus Wedekind

Department of Ecology and Evolution, Biophore, University of Lausanne, Lausanne, Switzerland

CW, 0000-0001-6143-4716

Major histocompatibility complex (MHC) social signalling has been found in over 20 vertebrate species so far and is 'likely the basis for a vertebrate-wide chemosensory communication system' [1]. Numerous further examples of MHC social signalling have been published since Ruff *et al.*'s [1] exhaustive review, both demonstrating female reactions to MHC sharing with males (e.g. [2,3]) and male reactions to MHC sharing with females (e.g. [4–7]). When concentrating on experimental studies in humans, 15 papers so far claimed to provide evidence for MHC-linked odours and/or odour preferences (electronic supplementary material, table S1), and a recent meta-analysis concluded that MHC-linked preferences are 'likely conserved across primates' [8]. Well-worked-out cases of absent MHC social signalling would therefore be interesting exceptions of what seems to be a general rule, and it is important to find and document such exceptions to learn more about the principles of social signalling. However, easy as it is to miss an existing effect (e.g. because of problematic experimental protocols or low statistical power), it is just as challenging to demonstrate that an effect does not exist.

Probst *et al.* [9] argue that they found an example of absent MHC social signalling. They studied men's preferences of women's body odours, following an experimental design that is largely analogous to the one Wedekind *et al.* [10] had used when they found women's preferences for men's odours to be MHC linked. Probst *et al.* [9] collected armpit odours from donors and presented eight of them to raters, with four of them being very MHC-dissimilar, and the other four very MHC-similar (i.e. testing only the extremes; Wedekind *et al.* [10] had presented three of the extremes each). Both groups studied Swiss students and tried to make sure that their samples were ethnically closely defined (ethnicity can be a confounding factor in studies of MHC social signalling [11,12]). However, there are important differences between the two studies that can lead to wrong conclusions.

First, if evidence for MHC social signalling is found [10], with preferences depending on MHC sharing, it is possible to conclude that there are effects of the MHC (or linked genes) on both sides of the communication (i.e. the donors' body odours contain MHC-linked signals and the raters' odour preferences are linked to their own MHC). If there were indeed no evidence for MHC social signalling, it remained unclear whether there is no influence of the MHC on the production of odours (donor effects) or on the perception of odours (rater effects), or on both. Probst *et al.* [9] proposed that the 'HLA has no effect on men's odour preferences' (their abstract; the human's MHC is called human leukocyte antigen, HLA). However, it is still possible that men have MHC-linked preferences and that, in Probst *et al.* [9], women's body odours did not contain MHC-linked signals, or that these signals were too weak to be detected. Wedekind & Füre [13] tested for MHC-linked odour preferences and found evidence for MHC-linked odour preferences in both men and women. They recorded the preferences of 121 male and female raters of the same six odours (from two female and four male donors) and found the amount of variance ( $r^2$ ) in

**Table 1.** Mixed-effects analyses of pleasantness scorings in Probst *et al.* [18] testing the effects of the sharing of MHC antigens between donor and rater (dissimilar versus similar; 'MHC') and odour intensity (intensity) as fixed factors, and rater identity or donor identity (ID) as random factors. The proportions of the total variance explained are based on REML variance component estimates (unbounded). Significant *p*-values are marked in italic.

fixed factors	pleasantness <sup>a</sup>		pleasantness <sup>b</sup>	
	<i>F</i>	<i>p</i>	<i>F</i>	<i>P</i>
MHC	0.4	0.55	1.5	0.23
Intensity	60.7	<0.001	35.0	<0.001
MHC × intensity	4.4	0.04	4.4	0.04
<i>random factors</i>				
ID (%)		4.4		7.4
ID × MHC (%)		0.6		0
ID × intensity (%)		43.8		0
ID × MHC × intensity (%)		17.8		51.8
residual (%)		33.5		40.8

<sup>a</sup>With rater identity as random factor.

<sup>b</sup>With donor identity as random factor.

pleasantness scoring that was explained by the sharing of MHC antigens between donors and raters to vary between nearly 0 and 22.6%. The highest  $r^2$  was found when male raters evaluated the odour of a male donor, while the  $r^2$  turned out to be low for the two female donors (i.e. the question of whether women signal their MHC was not solved yet; see also electronic supplementary material, table S1). A further consequence of the low number of odour donors in their study was that possibly confounding effects of non-MHC-linked stimuli could not be sufficiently excluded. However, evidence for MHC-linked preferences in both men and women was again found in another study on body odours [12], when studying preferences for traditional perfume ingredients [14], or in electroencephalograms of subjects smelling human odour samples (axillary hairs) [15] (electronic supplementary material, table S1). Male preferences for MHC-dissimilar females have also been found in other species (e.g. [4–7]). Given these repeated accounts of MHC-linked preferences in men or males, the most parsimonious explanation for the non-significant findings of Probst *et al.* [9] is that the odours of their donors did not contain strong MHC-linked signals. However, it would also be premature to conclude that there is no MHC social signalling in women. This is because of the second important deviation from previous studies.

In previous studies [10,13], all donors except one had unshaved armpits, while all donors in Probst *et al.* [9] were asked to shave their armpits shortly before collection of odours because 'axillary hair might affect the perceived quality of the axillary odour' (their electronic supplementary material S1). Indeed, odours of shaved armpits are typically perceived as less intense than odours of unshaved armpits [16]. Moreover, the density of apocrine glands is highest in the armpit and a few other regions, including the genitoanal regions, and the microbial processing of apocrine secretions that plays an important role in producing body odours is enhanced by axillary and pubic hair [17]. These may be some of the reasons why shaving armpits and using anti-microbial deodorants have become social norms among women in many parts of the world. If the functional significance of armpit hair is to enhance social signalling [15], shaving them reduces this type of signalling. We would then predict that

odour intensities and the statistical link to MHC would be reduced. Indeed, the raters in Probst *et al.* [9] stated in 37 cases that they 'cannot smell the sample' [9], while this never happened in the analogous study that allowed for axillary hairs [10]. Moreover, the odours that could be perceived were on average rated as less intense in Probst *et al.* [9,18] than in the analogous study that allowed for axillary hairs [10,19] (mean  $\pm$  s.e. =  $49.7 \pm 0.1$  versus  $60.5 \pm 0.1$  on a scale from 0 to 100;  $F_{1,1000} = 37.0$ ,  $p < 0.0001$ ). A reduced signal intensity could be responsible for the apparent absence of MHC social signalling [9]. However, Probst *et al.* [9] did not yet test all hypotheses about MHC social signalling that can be derived from previous studies.

Numerous psychophysical studies demonstrate that the relationship between mixtures of volatile chemicals and their perception can be complex and difficult to predict [20]. Empirical observations may therefore be required to obtain testable predictions. Wedekind *et al.* [10] observed that odour pleasantness correlated negatively with odour intensity if donors and raters were MHC dissimilar ( $p = 0.01$ ), but not if they were MHC similar ( $p = 0.98$ ; see their fig. 3). The causalities behind this link to MHC sharing is still unclear, but the observation [10] leads to the testable prediction that MHC social signals affect the perception of odour pleasantness in combination with odour intensity.

I used mixed models (in JMP; [www.jmp.com](http://www.jmp.com)) to test this hypothesis with the data provided by Probst *et al.* [18], predicting odour pleasantness by the two (fixed) factors odour intensity and MHC sharing (similar or dissimilar), while controlling for effects of rater identity or donor identity (random factors; controlling for donor identity is predicted to provide more statistical power than controlling for rater identity [21]). This reanalysis revealed an overall negative correlation between pleasantness and intensity of odours (as reported before [9]) that was, however, dependent on whether donors and raters were similar or dissimilar on their MHC (see significant interactions in table 1). As predicted [10], the link between pleasantness and intensity was significantly more pronounced in MHC-dissimilar pairs than in MHC-similar pairs (electronic supplementary material, figure S1). This finding seemed independent of whether the mixed model

controlled for rater or for donor identity (table 1). When combining the datasets of Probst *et al.* [18] and Wedekind *et al.* [10] in one statistical model, the effects of intensity and MHC sharing did not seem to change (electronic supplementary material, table S2). There were study differences in mean pleasantness scorings and interaction effects on pleasantness (electronic supplementary material, table S2) that tended to be stronger when controlling for donor identity than for rater identity. Importantly, however, the three-way interactions (study  $\times$  intensity  $\times$  MHC sharing) did not explain pleasantness (i.e. both studies found a similar intensity  $\times$  MHC interaction on pleasantness; electronic supplementary material, table S2).

In summary, Probst *et al.*'s [9] conclusions about MHC social signalling in humans were premature. Their experimental set-up would not allow to distinguish between an absence of MHC-linked odour preferences in men and an absence of MHC social signalling in women (while evidence for both

types of MHC effects had been found before in humans and other vertebrates). Moreover, their protocol reduced the chances of finding MHC-linked signals, because women were asked to shave their armpit hair. Nevertheless, a reanalysis of Probst *et al.*'s data [18] revealed a weak but statistically significant interaction between odour intensity and MHC sharing on odour pleasantness in the predicted direction [10]. The link between odour intensity and pleasantness was significantly stronger in MHC-dissimilar pairs than in MHC-similar pairs. Therefore, Probst *et al.*'s data [18] provide evidence for both MHC social signalling in women and MHC-linked odour preferences in men.

**Data accessibility.** This article has no additional data.

**Competing interests.** I declare I have no competing interests.

**Funding.** I received no funding for this study.

**Acknowledgements.** I thank I. Cuthill and the reviewers for comments and discussion.

## References

- Ruff JS, Nelson AC, Kubinak JL, Potts WK. 2012 MHC signaling during social communication. *Adv. Exp. Med. Biol.* **738**, 290–313. (doi:10.1007/978-1-4614-1680-7\_17)
- Santos PS *et al.* 2016 MHC-dependent mate choice is linked to a trace-amine-associated receptor gene in a mammal. *Sci. Rep.* **6**, 38490. (doi:10.1038/srep38490)
- Burger D, Thomas S, Aepli H, Dreyer M, Fabre G, Marti E, Sieme H, Robinson MR, Wedekind C. 2017 Major histocompatibility complex-linked social signalling affects female fertility. *Proc. R. Soc. B* **284**, 20171824. (doi:10.1098/rspb.2017.1824)
- Roth O, Sundin J, Berglund A, Rosenqvist G, Wegner KM. 2014 Male mate choice relies on major histocompatibility complex class I in a sex-role-reversed pipefish. *J. Evol. Biol.* **27**, 929–938. (doi:10.1111/jeb.12365)
- Burger D, Dolivo G, Marti E, Sieme H, Wedekind C. 2015 Female major histocompatibility complex type affects male testosterone levels and sperm number in the horse (*Equus caballus*). *Proc. R. Soc. B* **282**, 20150407. (doi:10.1098/rspb.2015.0407)
- Leclaire S, Strandh M, Mardon J, Westerdahl H, Bonadonna F. 2017 Odour-based discrimination of similarity at the major histocompatibility complex in birds. *Proc. R. Soc. B* **284**, 20162466. (doi:10.1098/rspb.2016.2466)
- Jeannerat E, Marti E, Berney C, Janett F, Bollwein H, Sieme H, Burger D, Wedekind C. 2018 Stallion semen quality depends on MHC matching to teaser mare. *Mol. Ecol.* **27**, 1025–1035. (doi:10.1111/mec.14490)
- Winternitz J, Abbate JL, Huchard E, Havlicek J, Garamszegi LZ. 2017 Patterns of MHC-dependent mate selection in humans and nonhuman primates: a meta-analysis. *Mol. Ecol.* **26**, 668–688. (doi:10.1111/mec.13920)
- Probst F, Fischbacher U, Lobmaier JS, Wirthmüller U, Knoch D. 2017 Men's preferences for women's body odours are not associated with human leucocyte antigen. *Proc. R. Soc. B* **284**, 20171830. (doi:10.1098/rspb.2017.1830)
- Wedekind C, Seebeck T, Bettens F, Paepke AJ. 1995 MHC-dependent mate preferences in humans. *Proc. R. Soc. Lond. B* **260**, 245–249. (doi:10.1098/rspb.1995.0087)
- Rosenberg LT, Cooperman D, Payne R. 1983 HLA and mate selection. *Immunogenetics* **17**, 89–93. (doi:10.1007/bf00364292)
- Thornhill R, Gangestad SW, Miller R, Scheyd G, McCollough JK, Franklin M. 2003 Major histocompatibility complex genes, symmetry, and body scent attractiveness in men and women. *Behav. Ecol.* **14**, 668–678. (doi:10.1093/beheco/arg043)
- Wedekind C, Furi S. 1997 Body odour preferences in men and women: do they aim for specific MHC combinations or simply heterozygosity? *Proc. R. Soc. Lond. B* **264**, 1471–1479. (doi:10.1098/rspb.1997.0204)
- Milinski M, Wedekind C. 2001 Evidence for MHC-correlated perfume preferences in humans. *Behav. Ecol.* **12**, 140–149. (doi:10.1093/beheco/12.2.140)
- Pause BM, Krauel K, Schraders C, Sojka B, Westphal E, Müller-Ruchholtz W, Ferstl R. 2006 The human brain is a detector of chemosensorily transmitted HLA-class I-similarity in same- and opposite-sex relations. *Proc. R. Soc. B* **273**, 471–478. (doi:10.1098/rspb.2005.3342)
- Kohoutova D, Rubesova A, Havlicek J. 2012 Shaving of axillary hair has only a transient effect on perceived body odor pleasantness. *Behav. Ecol. Sociobiol.* **66**, 569–581. (doi:10.1007/s00265-011-1305-0)
- Grice EA, Segre JA. 2011 The skin microbiome. *Nat. Rev. Microbiol.* **9**, 244–253. (doi:10.1038/nrmicro2537)
- Probst F, Fischbacher U, Lobmaier JS, Wirthmüller U, Knoch D. 2017 Data from: Men's preferences for women's body odours are not associated with HLA. Dryad Digital Repository. (doi:10.5061/dryad.270h8)
- Wedekind C, Seebeck T, Bettens F, Paepke AJ. 1995 Data from: MHC-dependent mate preferences in humans. Dryad Digital Repository. (doi:10.5061/dryad.r87pq)
- Ferreira V. 2012 Revisiting psychophysical work on the quantitative and qualitative odour properties of simple odour mixtures: a flavour chemistry view. Part 1. Intensity and detectability: a review. *Flavour Frag. J.* **27**, 124–140. (doi:10.1002/ffj.2090)
- Wedekind C, Seebeck T. 1996 MHC and mate selection in humans? Reply. *Trends Ecol. Evol.* **11**, 24–25. (doi:10.1016/0169-5347(96)80237-0)

*Electronic Supplementary Material 1*

**A predicted interaction between odour pleasantness and intensity provides evidence for MHC social signalling in women**

*C. Wedekind, Department of Ecology and Evolution, Biophore, University of Lausanne, Lausanne, Switzerland*

**Supplementary Table S1.** Studies that tested experimentally for MHC-linked body or urinary odours and/or odour preferences in humans, based on a search in Web of Science ([www.webofknowledge.com](http://www.webofknowledge.com)), starting with the keywords “HLA” and “odour”, and examining studies that cited, or were cited by, the papers that matched the search criteria. The table summarizes the conclusions drawn by the authors of the respective paper, with “yes” = the authors concluded that they had found evidence for MHC-linked effects on odours or odour perception; “(yes)” = both sexes were included and analyses were not sex-specific; “.“ = link to MHC was not tested; “?” = the role of the MHC remained unclear because there was no statistically significant link. The column “Critique & authors’ reply” refers to commentary papers that specifically addressed aspects of the respective paper.

Study	MHC-linked odours		MHC-linked odour perception		Comments	Critique & authors' reply
	men	women	men	women		
Gilbert et al. (1986) [1]	.	.	(yes)	(yes)	Humans evaluating odours of MHC-congenic mice.	
Ferstl et al. (1992) [2]	(yes)	(yes)	.	yes	Human urinary odours evaluated by rats; women reporting peculiar odour perception phenomena.	
Wedekind et al. (1995) [3]	yes	.	.	yes	Users of contraceptive Pill had different preferences than non-Pill users.	[4], [5]
Wedekind & Furi (1997) [6]	yes	?	yes	yes	No significant sex differences in MHC-linked preferences; strongest MHC-linked preference for one male odour ( $r^2 = 0.23$ ).	
Eggert et al. (1999) [7]	yes	yes	.	.	Human odours evaluated by rats and in gas chromatography.	
Milinski & Wedekind (2001) [8]	.	.	yes	yes	MHC-linked preferences for some traditional perfume ingredients.	
Jacob et al. (2002) [9]	yes	.	.	yes	Preferences tested in a group of low genetic diversity.	[10], [11]
Thornhill et al. (2003) [12]	yes	yes	yes	yes	Preferences of male odours linked to MHC heterozygosity, preferences of female odours linked to MHC similarity.	
Santos et al. (2005) [13]	yes	?	?	yes	Potential effects of the contraceptive pill not included in analyses.	
Pause et al. (2006) [14]	yes	yes	yes	yes	Analysing electroencephalograms with axillary hairs as odour samples.	
Wedekind et al. 2007 [15]	yes	.	?	yes	Descriptions of body odours by a female perfumer.	
Roberts et al. (2008) [16]	yes	.	.	yes	Odour preferences shifted towards MHC similarity with use of the contraceptive Pill (as predicted in [3])	
Janes et al. (2010) [17]	?	?	(yes)	(yes)	MHC-linked preferences to artificial scents, no significant link to body odours (the authors discuss ethnic diversity as potentially confounding).	
Natsch et al. (2010) [18]	?	?	.	.	Testing for MHC effects on N-acylglutamine conjugates of volatile carboxylic acids secreted in the axilla.	
Hämmerli et al. (2012) [19]	.	.	(yes)	(yes)	MHC-linked preferences for some traditional perfume ingredients	
Milinski et al. (2013) [20]	.	.	.	yes	Supplementation of own body odour by synthesized MHC peptides.	[21], [22]
Verhulst et al. (2013) [23]	?	.	.	.	Testing MHC effects on attractiveness of odours to mosquitoes.	
Probst et al. (2017) [24]	.	?	?	.		present paper
Present reanalysis of Probst et al.	.	yes	yes	.		

The table only lists studies that tested for potential MHC effects on odours or odour preferences in humans. It does not list studies that tested for links between MHC and facial or skin characteristics [e.g. 25, 26, 27] or that tested for potential MHC effects on mate choice, sexual responsivity, or fertility in humans for which some studies did not find significant MHC effects [e.g. 28, 29] while others found statistically significant links to the MHC [e.g. 30, 31, 32-34].

## References cited

1. Gilbert A.N., Yamazaki K., Beauchamp G.K., Thomas L. 1986 Olfactory discrimination of mouse strains (*Mus musculus*) and major histocompatibility complex types by humans (*Homo sapiens*). *Journal of Comparative Psychology* **100**(3), 262-265. (doi:10.1037//0735-7036.100.3.262).
2. Ferstl R., Eggert F., Westphal E., Zavazava N., Muller-Ruchholtz W. 1992 MHC-related odors in humans. In *Chemical Signals in Vertebrates VI* (eds. Doty R.L., Müller-Schwarze D.), pp. 205-211. New York, Plenum.
3. Wedekind C., Seebeck T., Bettens F., Paepke A.J. 1995 MHC-dependent mate preferences in humans. *Proc. R. Soc. B Biol. Sci.* **260**(1359), 245-249. (doi:10.1098/rspb.1995.0087).
4. Hedrick P.W., Loeschcke V. 1996 MHC and mate selection in humans? *Trends Ecol. Evol.* **11**(1), 24.
5. Wedekind C., Seebeck T. 1996 MHC and mate selection in humans? Reply. *Trends Ecol. Evol.* **11**(1), 24-25. (doi:10.1016/0169-5347(96)80237-0).
6. Wedekind C., Furi S. 1997 Body odour preferences in men and women: do they aim for specific MHC combinations or simply heterozygosity? *Proc. R. Soc. B Biol. Sci.* **264**(1387), 1471-1479. (doi:10.1098/rspb.1997.0204).
7. Eggert F., Luszyk D., Haberkorn K., Wobst B., Vostrowsky O., Westphal E., Bestmann H.J., Muller-Ruchholtz W., Ferstl R. 1999 The major histocompatibility complex and the chemosensory signalling of individuality in humans. *Genetica* **104**(3), 265-273. (doi:10.1023/a:1026431303879).
8. Milinski M., Wedekind C. 2001 Evidence for MHC-correlated perfume preferences in humans. *Behav. Ecol.* **12**(2), 140-149. (doi:10.1093/beheco/12.2.140).
9. Jacob S., McClintock M.K., Zelano B., Ober C. 2002 Paternally inherited HLA alleles are associated with women's choice of male odor. *Nat. Genet.* **30**, 175-179.
10. Wedekind C. 2002 The MHC and body odors: arbitrary effects caused by shifts of mean pleasantness. *Nat. Genet.* **31**, 237.
11. McClintock M.K., Schumm P., Jacob S., Zelano B., Ober C. 2002 The MHC and body odors: arbitrary effects caused by shifts of mean pleasantness - Reply. *Nat. Genet.* **31**(3), 237-238. (doi:10.1038/ng0702-237b).
12. Thornhill R., Gangestad S.W., Miller R., Scheyd G., McCollough J.K., Franklin M. 2003 Major histocompatibility complex genes, symmetry, and body scent attractiveness in men and women. *Behav. Ecol.* **14**(5), 668-678. (doi:10.1093/beheco/arg043).
13. Santos P.S.C., Schinemann J.A., Gabardo J., Bicalho M.D. 2005 New evidence that the MHC influences odor perception in humans: a study with 58 Southern Brazilian students. *Horm. Behav.* **47**(4), 384-388.
14. Pause B.M., Krauel K., Schraders C., Sojka B., Westphal E., Muller-Ruchholtz W., Ferstl R. 2006 The human brain is a detector of chemosensorily transmitted HLA-class I-similarity in same- and opposite-sex relations. *Proc. R. Soc. B Biol. Sci.* **273**(1585), 471-478. (doi:10.1098/rspb.2005.3342).
15. Wedekind C., Escher S., Van de Waal M., Frei E. 2007 The major histocompatibility complex and perfumers' descriptions of human body odors. *Evol. Psychol.* **5**, 330-343.
16. Roberts S.C., Gosling L.M., Carter V., Petrie M. 2008 MHC-correlated odour preferences in humans and the use of oral contraceptives. *Proc. R. Soc. B Biol. Sci.* **275**, 2715-2722.
17. Janes D., Klun I., Vidan-Jeras B., Jeras M., Kreft S. 2010 Influence of MHC on odour perception of 43 chemicals and body odour. *Central European Journal of Biology* **5**(3), 324-330. (doi:10.2478/s11535-010-0020-6).
18. Natsch A., Kuhn F., Tiercy J.M. 2010 Lack of evidence for HLA-linked patterns of odorous carboxylic acids released from glutamine conjugates secreted in the human axilla. *Journal of Chemical Ecology* **36**(8), 837-846. (doi:10.1007/s10886-010-9826-y).
19. Hämmerli A., Schweisgut C., Kaegi M. 2012 Population genetic segmentation of MHC-correlated perfume preferences. *International Journal of Cosmetic Science* **34**(2), 161-168. (doi:10.1111/j.1468-2494.2011.00696.x).


20. Milinski M., Croy I., Hummel T., Boehm T. 2013 Major histocompatibility complex peptide ligands as olfactory cues in human body odour assessment. *Proc. R. Soc. B Biol. Sci.* **280**(1755). (doi:10.1098/rspb.2012.2889).
21. Natsch A. 2014 A human chemosensory modality to detect peptides in the nose? *Proc. R. Soc. B Biol. Sci.* **281**(1776). (doi:10.1098/rspb.2013.1678).
22. Milinski M., Croy I., Hummel T., Boehm T. 2014 Reply to A human chemo-sensory modality to detect peptides in the nose? by A. Natsch. *Proc. R. Soc. B Biol. Sci.* **281**(1776). (doi:10.1098/rspb.2013.2816).
23. Verhulst N.O., Beijleveld H., Qiu Y.T., Maliepaard C., Verduyn W., Haasnoot G.W., Claas F.H.J., Mumm R., Bouwmeester H.J., Takken W., et al. 2013 Relation between HLA genes, human skin volatiles and attractiveness of humans to malaria mosquitoes. *Infection Genetics and Evolution* **18**, 87-93. (doi:10.1016/j.meegid.2013.05.009).
24. Probst F., Fischbacher U., Lobmaier J.S., Wirthmüller U., Knoch D. 2017 Men's preferences for women's body odours are not associated with human leucocyte antigen. *Proc. R. Soc. B Biol. Sci.* **284**, 20171830. (doi:10.1098/rspb.2017.1830).
25. Roberts S.C., Little A.C., Gosling L.M., Jones B.C., Perrett D.I., Carter V., Petrie M. 2005 MHC-assortative facial preferences in humans. *Biol. Lett.* **1**(4), 400-403.
26. Roberts S.C., Little A.C., Gosling L.M., Perrett D.I., Carter V., Jones B.C., Penton-Voak I., Petrie M. 2005 MHC-heterozygosity and human facial attractiveness. *Evol. Hum. Behav.* **26**(3), 213-226.
27. Lie H.C., Simmons L.W., Rhodes G. 2010 Genetic dissimilarity, genetic diversity, and mate preferences in humans. *Evol. Hum. Behav.* **31**(1), 48-58. (doi:10.1016/j.evolhumbehav.2009.07.001).
28. Hedrick P.W., Black F.L. 1997 HLA and mate selection: No evidence in South Amerindians. *Am. J. Hum. Genet.* **61**(3), 505-511.
29. Ihara Y., Aoki K., Tokunaga K., Takahashi K., Juji T. 2000 HLA and human mate choice: Tests on Japanese couples. *Anthropological Science* **108**(2), 199-214.
30. Ober C., Elias S., Kostyu D.D., Hauck W.W. 1992 Decreased fecundability in Hutterite couples sharing HLA-DR. *Am. J. Hum. Genet.* **50**(1), 6-14.
31. Ober C., Weitkamp L.R., Cox N., Dytch H., Kostyu D., Elias S. 1997 HLA and mate choice in humans. *Am. J. Hum. Genet.* **61**(3), 497-504.
32. Garver-Apgar C.E., Gangestad S.W., Thornhill R., Miller R.D., Olp J.J. 2006 Major histocompatibility complex alleles, sexual responsiveness, and unfaithfulness in romantic couples. *Psychological Science* **17**(10), 830-835. (doi:10.1111/j.1467-9280.2006.01789.x).
33. Kromer J., Hummel T., Pietrowski D., Giani A.S., Sauter J., Ehninger G., Schmidt A.H., Croy I. 2016 Influence of HLA on human partnership and sexual satisfaction. *Sci. Rep.* **6**, 32550. (doi:10.1038/srep32550).
34. Sapphire-Bernstein S., Larson C.M., Gildersleeve K.A., Fales M.R., Pillsworth E.G., Haselton M.G. 2017 Genetic compatibility in long-term intimate relationships: partner similarity at major histocompatibility complex (MHC) genes may reduce in-pair attraction. *Evol. Hum. Behav.* **38**, 190-196. (doi:10.1016/j.evolhumbehav.2016.09.003).

*Electronic Supplementary Material 2*

**A predicted interaction between odour pleasantness and intensity provides evidence for MHC social signalling in women**

*C. Wedekind, Department of Ecology and Evolution, Biophore, University of Lausanne, Lausanne, Switzerland*

**Supplementary Figure S1.** Pleasantness versus intensity of body odour of MHC-dissimilar (black symbols and line) and MHC-similar pairs (grey symbols and line) of odour donor and rater in Probst et al. [1]. The lines show the regressions. See Table 1 for statistics.



**Literature cited**

1. Probst F., Fischbacher U., Lobmaier J.S., Wirthmüller U., Knoch D. 2017 Data from: Men's preferences for women's body odours are not associated with HLA. In *Dryad Digital Repository* (10.5061/dryad.270h8)

### Electronic Supplementary Material 3

## A predicted interaction between odour pleasantness and intensity provides evidence for MHC social signalling in women

C. Wedekind, Department of Ecology and Evolution, Biophore, University of Lausanne, Lausanne, Switzerland

**Supplementary Table S2:** Mixed-effects analyses of pleasantness scorings in the combined datasets of Probst et al. [1] and Wedekind et al. [2], testing the effects of the sharing of MHC antigens between donor and rater (dissimilar versus similar; “MHC”), odour intensity (“intensity”), and “study” (Probst et al. [1] versus Wedekind et al. [2]) as fixed factors, and rater identity or donor identity (“ID”) as random factors. The proportions of the total variance explained are based on REML variance component estimates (unbounded). Significant p-values are marked in bold.

Fixed factors	Pleasantness <sup>1</sup>		Pleasantness <sup>2</sup>	
	F	p	F	p
MHC	2.0	0.16	1.0	0.33
Intensity	120.2	<b>&lt;0.001</b>	64.5	<b>&lt;0.001</b>
MHC x intensity	4.1	<b>0.05</b>	5.2	<b>0.02</b>
Study	35.0	<b>&lt;0.001</b>	15.2	<b>&lt;0.001</b>
Study x MHC	5.3	<b>0.02</b>	5.6	<b>0.02</b>
Study x intensity	2.2	0.14	5.3	<b>0.03</b>
Study x MHC x intensity	0.1	0.72	0.01	0.91
<i>Random factors</i>				
ID	0%		5.7%	
ID x MHC	1.9%		0%	
ID x intensity	0%		0%	
ID x MHC x intensity	54.9%		0%	
ID x study	2.9%		0%	
ID x study x MHC	0%		4.9%	
ID x study x intensity	37.3%		45.4%	
ID x study x MHC x intensity	0%		40.7%	
Residual	3.0%		3.3%	

<sup>1</sup> with rater identity as random factor; <sup>2</sup> with donor identity as random factor

### Literature cited

1. Probst F., Fischbacher U., Lobmaier J.S., Wirthmüller U., Knoch D. 2017 Data from: Men’s preferences for women’s body odours are not associated with HLA. In *Dryad Digital Repository* (10.5061/dryad.270h8)
2. Wedekind C., Seebeck T., Bettens F., Paepke A.J. 1995 Data from: MHC-dependent mate preferences in humans. In *Dryad Digital Repository* (10.5061/dryad.r87pq)